



Development and characterization of 108 SNP markers in the Iwagaki oyster, *Crassostrea nippona*

Kaikai Liu¹ · Hong Yu¹ · Qi Li^{1,2}

Received: 28 April 2018 / Accepted: 8 May 2018 / Published online: 10 May 2018
© Springer Science+Business Media B.V., part of Springer Nature 2018

Abstract

A total of 108 single nucleotide polymorphism (SNP) markers from the Iwagaki oyster, *Crassostrea nippona* were isolated and characterized based on high-throughput sequencing in this study. The minor allele frequency per locus ranged from 0.0167 to 0.4844. The observed heterozygosity and expected heterozygosity varied from 0 to 0.9688 and 0.0333 to 0.5079, respectively. Eighteen loci showed significant deviation from Hardy–Weinberg equilibrium. These SNP markers described in this study will provide a valuable tool for population genetic analysis and natural resource conservation of *C. nippona*.

Keywords *Crassostrea nippona* · SNP · High-throughput sequencing · High resolution melting

The Iwagaki oyster *Crassostrea nippona* is a commercially important bivalve, which is distributed along the coasts of Japan, Korea and China (Boudry et al. 2003; Itoh et al. 2004; Yoon et al. 2008). The commercial price of *C. nippona* is nearly five times as high as that of the Pacific oyster *C. gigas* in Japan, as it is edible during the summer when the other oyster species are unavailable (Itoh et al. 2004). Since the production of *C. nippona* is based almost exclusively on natural stocks, over-exploitation has led to continuing decline of the natural populations over the decades (Fujiwara 1995; Li 2007). Therefore, it is urgent to perform population genetic investigation on *C. nippona* to conserve the wild resources. Single nucleotide polymorphisms (SNPs) are now the most popular DNA markers in genetic studies because of their high level of polymorphism, favorable reproducibility and wide genomic distribution (Vignal et al. 2002). With the rapid development on next generation sequencing technologies, discovery of large numbers of SNPs in any non-model organisms of interest is becoming easier and faster. In this study, we developed and characterized SNP markers in *C. nippona* for the first time based on the restriction-site

associated DNA (RAD) sequencing. These markers will facilitate the researches on conservation genetics and genetic evaluation in *C. nippona*.

To obtain SNP marker resources, the RAD libraries were constructed and sequenced on the Illumina HiSeq 2500 platform using 150-bp paired-end reads. A total of 538,594 putative SNPs in *C. nippona* was identified, of which C/T and G/A were the most common substitution and the ratio of transition to transversion was 1.22. Four hundred and twenty putative SNPs were randomly chosen for primer design using Primer Premier 5.0 and evaluated for polymorphism by high resolution melting (HRM) analysis in a cultured population of *C. nippona* from Yantai, China (n=32). Genomic DNA was extracted from adductor muscle by standard proteinase K digestion and phenol–chloroform extraction. PCR was performed in a total volume of 10 µL on a Light Cycler[®] 480 real-time PCR instrument (Roche Diagnostics Burgess Hill, UK). The reaction mixture contained 0.25 U Taq DNA polymerase (Takara, Dalian, China), 10×PCR buffer, 0.2 mM dNTP mix, 0.4 µM of each primer set, 1.5 mM MgCl₂, 5 µM SYTO9 (Invitrogen Foster City, CA, USA), and about 10 ng template DNA. The PCR amplification procedure was as follows: first denaturation at 95 °C for 5 min, then 45–55 cycles of denaturation at 95 °C for 40 s, annealing and extension for 40 s at 62 °C for the first cycle and thereafter at 0.5 °C decrease each for 10 cycles, and a final extension at 72 °C for 40 s. Following amplification, the products were denatured at 95 °C for 1 min, and then annealed at 40 °C for 1 min to randomly form DNA duplexes. Melting curves were

✉ Hong Yu
hongyu@ouc.edu.cn

¹ Key Laboratory of Mariculture (Ocean University of China), Ministry of Education, Qingdao 266003, China

² Qingdao National Laboratory for Marine Science and Technology, Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao, China

Table 1 Characterization of 108 SNP markers in the Iwagaki oyster *Crassostrea nippona*

Locus ID	Primer sequence (5'–3')	Size (bp)	Tm (°C)	Types	Ho	He	MAF	F_{IS}	P_{HWE}
CNP1	F: TGTTGACGACTGAGAGCTGG R: CGATCACATACTGACGTGTCC	90	60	G/A	0.3548	0.2967	0.1774	– 0.2157	0.2553
CNP3	F: AGTGGTTTGAGGCACTCGTC R: TAGTTCTGGCCAAGCACCC	116	60	C/A	0.3667	0.3045	0.1833	– 0.2245	0.2442
CNP8	F: ATGCTGTTTCATGTTTGGA R: TCTGTTTGGGTTTCGGTTGT	113	60	G/T	0.3333	0.4520	0.3333	0.2500	0.1419
CNP10	F: ATTGCATAGCAAAGATCGCC R: ACATTCGGGATGAAGACTCG	124	60	C/A	0.1562	0.1463	0.0781	– 0.0847	0.6704
CNP11	F: AATTCGCCTTATTTACTTAGAGGA R: ATCTCTGAAACACCATCCGC	113	60	G/C	0.0645	0.4823	0.3871	0.8640	0.0000*
CNP14	F: ATAAGTCACATGCGCGGTTTC R: GAAAAAGCACACCAAAGCGA	85	60	A/C	0.0000	0.4528	0.3333	1.0000	0.0000*
CNP16	F: CGCAAGATTTGACAATTTCTGT R: CCACTTCGATTTCAATTTGTG	111	60	G/A	0.3548	0.2967	0.1774	– 0.2157	0.2553
CNP19	F: CTCCGCCCTTTAAAAGTCT R: AAAAGTCTGAAATTGCTTGCG	124	60	C/T	0.1250	0.1190	0.0625	– 0.0667	0.7458
CNP24	F: TGGCCATTGGTAAGTAAGGC R: AAGCGAGCATCTGTCTGATCT	115	60	C/T	0.0000	0.3913	0.2593	1.0000	0.0000*
CNP26	F: CGGCCAAGAATACCTTGACT R: CTACTTGGATGCTTCCGGTG	83	60	G/A	0.2500	0.2222	0.1250	– 0.1429	0.4528
CNP27	F: CCCATTGCACAACGTTACAA R: CCGGGGTTACGTGGTATTTA	104	60	T/G	0.3571	0.2987	0.1786	– 0.2174	0.2786
CNP38	F: GGAAGAAAACCAAGTGCATCAA R: ATGAACTTTTTGGCGACCTG	86	60	C/G	0.0312	0.0908	0.0469	0.6503	0.0000*
CNP41	F: TCGTGTCTTTCTTCGCCTTT R: AAGGCACTCTGAAACAAGCC	102	60	T/G	0.1250	0.1190	0.0625	– 0.0667	0.7458
CNP45	F: TCTAGGCCATTTATCCGTGG R: CAGAAAATGCACTCCCTTGG	87	60	C/T	0.0938	0.0908	0.0469	– 0.0492	0.8216
CNP55	F: TGCAGGAAGAGGATGATCTG R: ATAAGCAACAAACCGCAACC	113	60	A/G	0.7097	0.4654	0.3548	– 0.5500	0.0029
CNP67	F: TTTGAGAGAATTGCAGCCG R: CAGTCTGGATTTCTCTGCTGG	82	60	T/C	0.0645	0.0635	0.0323	– 0.0333	0.8964
CNP68	F: CCACTTCACTTTTTGGGACA R: CCCCCAATTTCCTAAAGTCC	91	60	C/T	0.3871	0.3173	0.1935	– 0.2400	0.2036
CNP79	F: TGCTTTGACCTTTCAATATACA R: TGCTCTAACGGCCTTTTGT	99	60	C/G	0.0625	0.0615	0.0312	– 0.0323	0.8981
CNP91	F: TGTCCGCCCTTCTTACAAAC R: CAAATAAACGCACCCCACTT	103	60	G/A	0.5938	0.4241	0.2969	– 0.4222	0.0207
CNP95	F: AGTTCATCCCGTATTGCGTC R: ACAATGCTTCAAAACTCGCC	117	60	G/A	0.2333	0.2593	0.1500	0.0850	0.5632
CNP105	F: GCAGTCTCTGAAGAGCAGGAA R: CACGTTTTTGCCATGCTGTA	98	60	C/T	0.5938	0.4955	0.4219	– 0.2172	0.2543
CNP113	F: AGGACATTTCTGTTTCGAT R: CCATCTTTTGGGATACAGGC	99	60	G/A	0.0645	0.0635	0.0323	– 0.0333	0.8964
CNP124	F: AAGGCGCTGTAACCAAGAG R: GATGAAGAGAGGCGTTTTGC	104	60	T/C	0.3333	0.2825	0.1667	– 0.2000	0.3020
CNP133	F: CGTAGTCTTTGGTTTGCAATCTT R: CAGGGAAGGAACTGCAAGAG	89	60	G/A	0.2903	0.2522	0.1452	– 0.1698	0.3020
CNP137	F: GCAGCTGAAATGTTCTCCT R: GGAAATAGAGACCAAATACGCA	117	60	T/C	0.5938	0.4241	0.2969	– 0.4222	0.0207
CNP145	F: CCTCTTTCGAGAAAATAACCCA R: TCAAAGCAGAGCAACAAGGA	115	60	G/T	0.2069	0.1887	0.1034	– 0.1154	0.5736
CNP150	F: AAGAAGCACACATACCGCCT R: AACTGTTCCGGTGAAGCCAG	118	60	T/G	0.0000	0.4950	0.4194	1.0000	0.0000*

Table 1 (continued)

Locus ID	Primer sequence (5'–3')	Size (bp)	T _m (°C)	Types	H _o	H _e	MAF	F _{IS}	P _{HWE}
CNP151	F: GGGTGTACCTTTTCCACAA R: GTGCCTTGGTCCTTACCAG	82	60	T/C	0.4138	0.3339	0.2069	–0.2609	0.1815
CNP155	F: CAGTTTGAACCTTTACGCCC R: TTAGCCGGAGGAAATAAAAGG	107	60	G/T	0.5484	0.4045	0.2742	–0.3778	0.0423
CNP165	F: CTTCTGGGGAACAGCAACAT R: GGTGTAAGCAGGTTTAGCCG	90	60	A/G	0.2581	0.2285	0.1290	–0.1481	0.4436
CNP166	F: TTCCAAGTCAACGCAGTTTG R: TGAACGGTTTGTCTTGCATC	87	60	C/A	0.2857	0.2494	0.1429	–0.1667	0.4130
CNP170	F: GGCTGTTTTTCTCGAAGCATT R: CTATGACGCTCCACAGCCAC	85	60	G/A	0.9688	0.5074	0.4844	–0.9394	0.0000*
CNP171	F: AATCAGATTCTGGGCTTTGC R: AACCTTCGACCTTTGACTG	111	60	G/A	0.8750	0.5000	0.4375	–0.7778	0.0000*
CNP176	F: ATGGAGAAGCAATTTGGGC R: CCCAGTTCGAGAAATACCA	98	60	A/G	0.0625	0.0615	0.0312	–0.0323	0.8981
CNP178	F: GCCTCCTAAACTCCAAATTTCC R: GCTGATAGAGAACCAAATGTTGA	93	60	T/C	0.7742	0.4823	0.3871	–0.6316	0.0006
CNP180	F: CGAACAATGTTTTAGCCCGA R: TGGTACTGTGGTGTGCGAAG	101	60	T/C	0.0000	0.4881	0.4000	1.0000	0.0000*
CNP182	F: ATAACCAGCTGCATTTTGCC R: CGAAGAACAAGACGACCTGG	101	60	T/C	0.0000	0.4791	0.3793	1.0000	0.0000*
CNP189	F: TGCTTCAGATTAATGCCGTTT R: ATCGGAAAACGCGTTACAAA	106	60	T/C	0.0667	0.0655	0.0333	–0.0345	0.8946
CNP192	F: TGCTACCATTTTCTCAGCC R: TCATCAAGCGTGCCTTAGTG	90	60	C/T	0.9688	0.5074	0.4844	–0.9394	0.0000*
CNP196	F: ACCCTGAACATCAAAATGCC R: GCGTAAAGAAGAAACCAGCG	120	60	G/T	0.8387	0.4950	0.4194	–0.7222	0.0001*
CNP197	F: AGACAGACTCGCTTTCAGCC R: CCTCCTCTGGTTCGTCTGTG	90	60	A/G	0.8966	0.5033	0.4483	–0.8125	0.0000*
CNP199	F: ATAGGCTGACGCTGATTGGT R: GGTTCCTTCGTGTTGACGTG	115	60	G/A	0.6250	0.4365	0.3125	–0.4545	0.0126
CNP200	F: TTCAGGAGGTCTGATACCCAA R: GACTTCGATTTGCACCTTCC	101	60	C/T	0.7500	0.4762	0.3750	–0.6000	0.0009
CNP202	F: TGCCGTTTCATGACTTACGTG R: ATCGGAAAACGCGTTACAAA	93	60	G/T	0.0625	0.0615	0.0312	–0.0323	0.8981
CNP204	F: TGATAAACCTCTGCTCGCAA R: AAGTGGCATGGGTCTAGGAG	85	60	A/G	0.5667	0.5079	0.4833	–0.1346	0.5193
CNP209	F: AGGGGCCTATCTGCATTTCT R: ATTTTCCGTGAAAGGGTGTG	98	60	G/A	0.0000	0.1190	0.0625	1.0000	0.0000*
CNP225	F: AATCAGCTTTGATTCTGTGGC R: AAGCAACAACAACACAGAGGAA	91	60	G/A	0.3438	0.2892	0.1719	–0.2075	0.2660
CNP233	F: CTGGCTTCAATCAGGTCACA R: GCAAGTGCAAGCTTTCCAAC	95	60	C/T	0.9688	0.5074	0.4844	–0.9394	0.0000*
CNP242	F: CAAGTGCCAATGTAACCCCT R: TGTTGACAGATGTCAAAAGC	89	60	A/G	0.1250	0.1190	0.0625	–0.0667	0.7458
CNP252	F: AAAGGGATGCAACTCTTGGA R: TGACTCAATACATGCCAGAACA	92	60	T/C	0.4516	0.3554	0.2258	–0.2917	0.1198
CNP254	F: GATTGCTGACGGTGTGTTGTG R: AAAACCTCAAGGTGGATTGAGA	123	60	A/G	0.0000	0.1831	0.1000	1.0000	0.0000*
CNP256	F: TTTTCATGGTAGATGAGTAGCATCC R: GCGAGGAAATCCGAGTCTTA	86	60	C/T	0.7407	0.4752	0.3704	–0.5882	0.0030
CNP259	F: AGTCAGCCCTGGAGCACTTA R: GTGAAGCCAGTCTTGAAGC	116	60	T/C	0.2258	0.2036	0.1129	–0.1273	0.5148
CNP269	F: TGGAATGTTTCATTTCCGC R: TGTCAGCAAAGTGTACAAAAAGG	115	60	A/C	0.3448	0.4065	0.2759	0.1369	0.4011

Table 1 (continued)

Locus ID	Primer sequence (5'–3')	Size (bp)	T _m (°C)	Types	<i>H_o</i>	<i>H_e</i>	MAF	<i>F_{IS}</i>	<i>P_{HWE}</i>
CNP270	F: ATCCCATGGTGCATTCAAGT R: CATAGGAGGACTCGGCTGAC	81	60	G/T	0.3750	0.3095	0.1875	– 0.2308	0.2142
CNP273	F: TGGAGCGCATTTCATATAA R: TGCCAATTGTAATGGACGAT	91	60	C/T	0.2812	0.2455	0.1406	– 0.1636	0.3860
CNP274	F: ACCCATTAGAGGTCGAGGGT R: GAGCATTTTAACACCCGTGC	105	60	T/C	0.2812	0.2455	0.1406	– 0.1636	0.3860
CNP282	F: TGGCAAACCTGTTCGGTATCC R: TCCTCATCGCTTACATTCCA	97	60	C/T	0.7143	0.4675	0.3571	– 0.5556	0.0043
CNP283	F: ACCGGTAATTTGAACACCGA R: GGGGGTTGTTTACTTAGGTGC	98	60	C/T	0.5556	0.4088	0.2778	– 0.3846	0.0549
CNP288	F: GGGGTTCCGTTGGAATTATC R: TGTGGGAGTACCTTTTGGC	85	60	G/A	0.4194	0.3728	0.2419	– 0.1433	0.4744
CNP291	F: GGGGGAACACCTGTCACTAA R: CCCTCCAGAATCAGACATCC	112	60	T/C	0.6452	0.4442	0.3226	– 0.4762	0.0101
CNP292	F: GTTCAACGAGCACCTTCTC R: TTTACCTGGAAAGACCCTGC	84	60	A/C	0.3125	0.2679	0.1562	– 0.1852	0.3235
CNP294	F: CTCATGCCTTTGGAATGGTT R: TCGGTGTTTACTTTTTGCATCTT	92	60	T/C	0.1935	0.1777	0.0968	– 0.1071	0.5888
CNP295	F: TCCTGTGTACAGATAAAGCTCCA R: GACATTCACAGATACACAGCCC	117	60	T/C	0.2812	0.2455	0.1406	– 0.1636	0.3860
CNP298	F: GCAAAATTCAGTGGTAGAGGAAA R: CAGATCCCTGTGTATAAGGACCA	81	60	G/A	0.4828	0.3727	0.2414	– 0.3182	0.1008
CNP299	F: TCAAGGCAAAATGGATTCTATG R: GGATGGTTTTGTATGCCGAC	86	60	C/T	0.2143	0.1948	0.1071	– 0.1200	0.5653
CNP303	F: TTCAAAGAATCGCCATAGCA R: CTTTCAAGATTTCCGGATGGAG	107	60	G/A	0.3548	0.2967	0.1774	– 0.2157	0.2553
CNP307	F: TTGCCCCCTAAATGAACATC R: AATTGTGGGCATTTGGATCA	110	60	A/G	0.1875	0.1726	0.0938	– 0.1034	0.5958
CNP308	F: CTATCCAGGAGCCTTTGTGC R: CAGGCAATGAAGGGGACTTA	116	60	C/T	0.6429	0.4442	0.3214	– 0.4737	0.0153
CNP312	F: CTCTCTCCAATGGAAAAACAGA R: GTTCCGGAATCCTTTTGGT	80	60	T/C	0.7500	0.4762	0.3750	– 0.6000	0.0009
CNP315	F: CTGTAAGCGATTTCGATCGTG R: GAAACCGATCGGAGTTCAAA	85	60	G/A	0.3125	0.2679	0.1562	– 0.1852	0.3235
CNP316	F: GAGGATGCATTATCAGGGGA R: TTGACAATGAAAGTGTGTGGC	83	60	A/T	0.1379	0.1307	0.0690	– 0.0741	0.7319
CNP319	F: AATTCCTGTTCGGATACCCAG R: ACAAACTAGCAGCGGAGGA	92	60	G/A	0.5161	0.3892	0.2581	– 0.3478	0.0622
CNP320	F: AATTCACTTGATTGGCATCC R: GCTTCGAGAAAAATGGTTGG	126	60	C/T	0.0625	0.0615	0.0312	– 0.0323	0.8981
CNP321	F: ATCTCGTCGTCGATGGAATC R: TTTTGGGAATTGTGGGGTA	94	60	T/G	0.6875	0.4583	0.3438	– 0.5238	0.0039
CNP322	F: TCCAAAGTTTGTCAATGCTGA R: AGAAAACCTGTTCCAATGCG	97	60	G/C	0.3667	0.3045	0.1833	– 0.2245	0.2442
CNP324	F: TCCCAAGCCAACTCCTAAA R: AATTGGACATGTGGGTCTCTC	93	60	C/T	0.4062	0.3646	0.2344	– 0.1320	0.5057
CNP326	F: ATATTTGAGGAAACGGGGC R: GTTGCCATTAACGGCTGTA	118	60	T/C	0.4375	0.3472	0.2188	– 0.2800	0.1291
CNP331	F: GAACCTGCCACCAACGAAAT R: TGTTCTGTGGTCATTTTCTCTG	82	60	T/C	0.5000	0.3814	0.2500	– 0.3333	0.0795
CNP333	F: GCCAATGTTACACCAACACG R: AGAGCAAACACACCTGAGGG	120	60	G/A	0.4688	0.3646	0.2344	– 0.3061	0.0962
CNP341	F: GATGACCCGGTAGTTGTGCT R: GTCGTAAGGGGGATGGGATA	86	60	A/C	0.4516	0.3554	0.2258	– 0.2917	0.1198

Table 1 (continued)

Locus ID	Primer sequence (5'–3')	Size (bp)	T _m (°C)	Types	H _o	H _e	MAF	F _{IS}	P _{HWE}
CNP344	F: CGTCAACATATTGCTGGCTG R: TGCACAGGATTTTAAAGACGG	84	60	A/G	0.8387	0.4950	0.4194	–0.7222	0.0001*
CNP347	F: TCCCTCTTGCTCCAGCTCTA R: CACATTCATGGTCAAGGCAC	124	60	A/G	0.7097	0.4654	0.3548	–0.5500	0.0029
CNP352	F: GGCATTGTTTAAAACCTCGTG R: GTGCGATCTCCGGCTAAATA	119	60	C/A	0.0333	0.0333	0.0167	–0.0169	1.0000
CNP353	F: AGCATTGTGTTTCTCCTCC R: TGGCAGTGACCAGTATGTGTG	104	60	G/A	0.0333	0.4944	0.4167	0.9314	0.0000*
CNP356	F: CTTTCCCCTTTGTCACTACT R: TGTAGGAAGCTGGCAGTGAA	91	60	T/C	0.1290	0.3173	0.1935	0.5867	0.0006
CNP357	F: AGGAAACGTCGCAACTCAAC R: GATTTGATAGGGCCCTTGGT	96	60	G/C	0.1875	0.1726	0.0938	–0.1034	0.5958
CNP365	F: CAATAGCAAGCTGTTGGTGC R: CCAATGCCTGATTGTCATTCT	92	60	T/C	0.5172	0.3902	0.2586	–0.3488	0.0712
CNP368	F: ACAGTTTGTGTCTTCATCACGG R: CCCACAATTCTGTGCTGCTA	80	60	A/G	0.4815	0.3725	0.2407	–0.3171	0.1158
CNP372	F: CACATATGGCTAAGACCCCG R: TCAAAAATGACTTAATCCTGTCCA	91	60	A/G	0.4828	0.3727	0.2414	–0.3182	0.1008
CNP375	F: ACGCGTCATCTGCAATCATA R: TGGTTTTGCCTTTTAAAGTACGA	81	60	G/A	0.1290	0.1227	0.0645	–0.0690	0.7414
CNP377	F: GCTTCGGTCTAAGTCTCCG R: ATGGCTTTGTGGTAACCGAG	81	60	A/C	0.6897	0.4598	0.3448	–0.5263	0.0059
CNP379	F: GCAAAGTTGAAGAAAAGAACTCC R: GATCTGGTCTTGGTTGGGAA	98	60	C/T	0.1562	0.2892	0.1719	0.4511	0.0067
CNP381	F: TTCGTTGTACAGACAAGCAACA R: CCCTGAACAGGTGTGTCAAA	91	60	T/C	0.4000	0.3254	0.2000	–0.2500	0.1927
CNP382	F: TGCTAGCTGTTGTCACTCGG R: TGGTGTACCTGACAGTCCCT	85	60	T/C	0.1379	0.1307	0.0690	–0.0741	0.7319
CNP384	F: CGCATAATGATGGCGATTCT R: GTTCTCTCCCTGTCAACCCAA	109	60	A/G	0.0625	0.0615	0.0312	–0.0323	0.8981
CNP385	F: ACCCAAACTACGAGGACG R: TCCTTCATAGCTCGTTACTGACC	85	60	T/C	0.4062	0.3289	0.2031	–0.2549	0.1685
CNP386	F: CCTCGCAAGAAACTACGCTT R: AGCAGCCAGTTGAAGTGTT	117	60	A/G	0.5806	0.4188	0.2903	–0.4091	0.0276
CNP387	F: TCATCTTGGAGCCTCAGTTG R: CTGCCATTCATCAACTGCTC	80	60	A/G	0.3125	0.2679	0.1562	–0.1852	0.3235
CNP389	F: CGTGCATGATAGCATAATTCC R: GCGGGCAGATCGATTAGTAT	110	60	G/T	0.6786	0.4565	0.3393	–0.5135	0.0085
CNP390	F: CAGTCGAAGACAAATGGCAA R: GAAAATTGTGTACCTTCCGCA	102	60	T/C	0.1562	0.1463	0.0781	–0.0847	0.6704
CNP391	F: AAGTGCATCAATTTCTGTGGA R: CAGAGCCAGCTTGTGATTT	106	60	G/A	0.1667	0.1554	0.0833	–0.0909	0.6586
CNP400	F: ATACTCCGACGCCAAAGATG R: TAAGGGACTGTTCTCGGCAT	96	60	C/G	0.0938	0.0908	0.0469	–0.0492	0.8216
CNP402	F: ATTCTGTCCGCTTTTGGAC R: CGATCAACATTGCCTCTTCA	114	60	G/T	0.0625	0.0615	0.0312	–0.0323	0.8981
CNP405	F: TCTTACCCATCTCCATAGAAA R: AGGCAACATTTGCTAAAGCC	91	60	A/G	0.2500	0.2222	0.1250	–0.1429	0.4528
CNP409	F: CCTTTGGGTGGAGAAAACAA R: GCCTTGTCCATGTGGAATCT	116	60	G/A	0.8387	0.4950	0.4194	–0.7222	0.0001*
CNP417	F: TCTTTAAAAGCCCTCCCT R: GCGGATACTAATTCCTTGCG	101	60	T/G	0.0938	0.0908	0.0469	–0.0492	0.8216
CNP420	F: GATGCCTGCCTTCAATCAAT R: AAATTCTTTTCCCTCTCCAGC	81	60	C/T	0.4815	0.3725	0.2407	–0.3171	0.1158

H_o observed heterozygosity, H_e expected heterozygosity, MAF minor allele frequency, F_{IS} inbreeding coefficient, P_{HWE}, the P values for Hardy–Weinberg equilibrium test (*P < 0.05/108 = 0.0005)

generated by heating samples from 60 to 90 °C with 25 data acquisitions per degree. At the end of each PCR reaction, the Light Cycler® 480 Gene Scanning Software was used to determine genotypes by analyzing the peaks in the melting curve. The minor allele frequency (MAF), expected heterozygosity (H_e), observed heterozygosity (H_o), inbreeding coefficient (F_{IS} ; Weir and Cockerham 1984) and χ^2 tests of deviations from Hardy–Weinberg equilibrium (HWE) were calculated using Popgene 1.32 (Yeh et al. 2000).

Of the 420 primer pairs, 108 SNP loci (25%) were polymorphic and produced distinct melting curves that can be genotyped by HRM (Table 1). The minor allele frequency was detected to be from 0.0167 to 0.4844. The observed heterozygosity (H_o) and expected heterozygosity (H_e) varied from 0 to 0.9688 and 0.0333 to 0.5079, respectively. The values of F_{IS} were estimated from -0.9394 to 1.0000 . Eighteen SNPs showed significant deviation from HWE after the Bonferroni correction ($P < 0.0005$). These polymorphic SNP markers will be useful for further population genetic analysis, natural resource conservation and selective breeding of *C. nippona*.

Acknowledgements This work was supported by Shandong Province (2014GHY115002), Fundamental Research Funds for the Central Universities (201762014), and Industrial Development Project of Qingdao City (17-3-3-64-nsh).

References

- Boudry P, Heurtebise S, Lapègue S (2003) Mitochondrial and nuclear DNA sequence variation of presumed *Crassostrea gigas* and *Crassostrea angulata* specimens: a new oyster species in Hong Kong? *Aquaculture* 228(1–4):15–25
- Fujiwara M (1995) The problems in seed production of Iwa oyster *Crassostrea nippona*. *Bull Kyoto Inst Ocean Fish Sci* 18:14–21 (in Japanese)
- Itoh N, Tun KL, Komiyama H, Ueki N, Ogawa K (2004) An ovarian infection in the Iwagaki oyster, *Crassostrea nippona*, with the protozoan parasite *Marteilioides chungmuensis*. *J Fish Dis* 27:311–314
- Li WJ (2007) Biology and cultivation of oyster *Crassostrea nippona*. *Fisheries Sci* 26:689–690 (in Chinese)
- Vignal A, Milan D, SanCristobal M, Eggen A (2002) A review on SNP and other types of molecular markers and their use in animal genetics. *Genet Sel Evol* 34:275–305
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evol Int J Org Evol* 38:1358–1370
- Yeh FC, Yang R, Boyle T, Ye Z, Xiyun J (2000) Popgene 1.32: Microsoft Windows-based freeware for population genetic analysis. Molecular Biology and Biotechnology Centre, University of Alberta, Edmonton, Alberta
- Yoon HS, Jung HT, Choi SD (2008) Suminoe oyster culture in Korea. *J Shellfish Res* 27:505–508